

On the Role of Locality Condition in Bell's Theorem

Habibollah Razmi

*Department of Physics, Faculty of Sciences, University of Qom , P.O.Box
37135-147, Qom, I.R. Iran.
razmi@qom.ac.ir*

Abstract

For a special stochastic realistic model in certain spin-correlation experiments and without imposing the locality condition, an inequality is found. Then, it is shown that quantum theory is able (is possible) to violate this inequality. This shows that, irrelevance of the locality condition, the quantum entanglement of the spin singlet-state is the reason for the violation of Bell's inequality in Bell's theorem.

PACS number: 03.65.Ud

Locality condition has an important role in many models of Bell's theorem. In Bell's original work [1], CH model [2], and almost all the other versions, a condition often used to derive different forms of Bell's inequality is the so-called locality condition. In fact, in many versions of Bell's theorem, in addition to realism (hidden variable(s)), at least, the locality condition is considered as a necessary assumption. Although in stochastic realistic models (e.g. CH model) the condition of determinism is not present, the so called locality condition is still one of the necessary assumptions. Here, we want to show that even before introducing the locality condition, quantum theory is in conflict with realistic stochastic models.

Consider a source which emits a pair of spin $\frac{1}{2}$ identical particles in a singlet-state. The two particles of the pair travel in opposite directions towards suitable measuring devices D_A and D_B . Identify each pair of escaping particles by an index i ($i = 1, 2, \dots, n$). The experimental arrangement is such that one particle from each pair will enter D_A , and the other one will enter D_B . Each detector has a preferred direction. The azimuthal orientation of this direction denoted by Q_A and Q_B according to whether one deals with θ_A or θ_B . Denote by r_{ji} the result obtained by D_j in the i th pair. This can take values $+1$ or -1 according to whether the deflection is along the preferred direction or its opposite. Each detector can take two possible values θ'_j and θ''_j . A crucial requirement is that the experimental set-up in sides A and B be spatially separated (i.e. having space-like separation in the sense of the special theory of relativity). There are four possible experimental specifications corresponding to the four possible pairs (θ'_A, θ'_B) , (θ'_A, θ''_B) , (θ''_A, θ'_B) and (θ''_A, θ''_B) . By definition the correlation function $C(r_A, r_B)$ is the average of the product of the results obtained by D_A and D_B :

$$C(r_A, r_B) = \frac{1}{n} \sum_{i=1}^n r_{Ai}(\theta_A, \theta_B) r_{Bi}(\theta_A, \theta_B) \quad (1)$$

Usually, the following locality condition is imposed [3-4]:

$$\begin{cases} r_A(\theta_A, \theta_B) = r_A(\theta_A) \\ r_B(\theta_A, \theta_B) = r_B(\theta_B) \end{cases} \quad (2)$$

and the correlation function (1) is written as:

$$C(r_A, r_B) = \frac{1}{n} \sum_{i=1}^n r_{Ai}(\theta_A) r_{Bi}(\theta_B) \quad (3)$$

and then an inequality (e.g. Bell's inequality) is found which cannot be always satisfied by quantum mechanics (Bell's theorem).

Here we are going to show that even before the introduction of the locality condition (2) (i.e. by using only the correlation function (1)), there is a discrepancy between the model used and quantum theory (Bell's theorem without locality condition).

Since both r_{Ai} and r_{Bi} are dichotomic variables (i.e. take values +1 or -1 only), their product is also dichotomic. Assume that from n runs of the experiment, m of them show the value +1 for the product $r_A r_B$; thus, $n - m$ of them would show the value -1. Therefore, the correlation function (3) is reduced to the following form:

$$C(r_A, r_B) = \frac{1}{n} ([\sum_{i=+1}^m (+1)] + [\sum_{i=m+1}^n (-1)]) \quad (4)$$

Or:

$$C(r_A, r_B) = \frac{2m}{n} - 1 \quad (5)$$

Here, we should mention that although m has a stochastic and unknown value, it is an integer between 0 and n depending on the experimental arrangement (orientation) of the devices.

For Future use, specially for the simplicity of algebra, let us define the following S -function as:

$$S(r_A, r_B) = \frac{[1 + C(r_A, r_B)]}{2} \quad (6)$$

which by means of (5), takes the form:

$$S(r_A, r_B) = \frac{m}{n} \quad (7)$$

One may consider N experminets all with n runs and similar in set-up but with arbitrary direction of detectors. Then, for each of these experiments the S -function is:

$$S_l = \frac{m_l}{n} \quad l = 1, 2, \dots, N \quad (8)$$

If in all these experiments the special "deterministic" case which corresponds to $\theta_A^l = \theta_B^l$ is excluded then all m_l s satisfy:

$$0 < m_l \leq n \quad (9)$$

This is because the special case $m_l = 0$ only corresponds to setting $\theta_A^l = \theta_B^l$.

We are interested to the following Γ -function corresponding to the results of the N experiments:

$$\Gamma = \sum_{l=1}^N S_l = \sum_{l=1}^N \frac{m_l}{n} \quad (10)$$

A comparison of (9) and (10) results in:

$$\Gamma \geq \frac{N}{n} \quad (11)$$

This inequality may be considered as a special kind of "Bell's inequality" for the stochastic realistic model considered here (with N experiments each with n runs and with arbitrary direction of detection excluding the special "deterministic" case with $\theta_A^l = \theta_B^l$).

It is important to emphasize once again that the inequality (11) has been derived without imposing the locality condition (2).

Now, the question is: Does quantum theory satisfy the inequality (11)?

The quantum mechanical result for the correlation function of the above-mentioned experimental set-up (singlet-state) is [5]:

$$C_{QM}(r_A, r_B) = -\cos(\theta_A - \theta_B) \quad (12)$$

Thus:

$$S_{QM}(r_A, r_B) = \sin^2\left(\frac{\theta_A - \theta_B}{2}\right) = \sin^2\left(\frac{\theta_{AB}}{2}\right) \quad (13)$$

where $\theta_{AB} = \theta_A - \theta_B$.

Therefore:

$$\Gamma_{QM} = \sum_{l=1}^N \sin^2\left(\frac{\theta_{AB}^l}{2}\right) \quad (14)$$

This Γ_{QM} is able (is possible) to violate the inequality (11). Let show this in more detail.

Suppose all θ_{AB}^l s are selected so that:

$$0 < \theta_{AB}^l < 2\text{Arcsin}\left(\frac{1}{\sqrt{n}}\right) \quad l = 1, 2, \dots, N \quad (15)$$

Then:

$$\Gamma_{QM} < \frac{N}{n} \quad (16)$$

This is in conflict with the inequality (11). It means that one can always find cases for them quantum theory cannot reproduce the predictions of the special stochastic realistic (not necessarily local) model. This can be considered as a model of Bell's theorem without locality condition.

In fact, in the correlation function introduced in (1), the factorization of the terms under sum in the form of $r_A \cdot r_B$, irrelevance of the functional form of r_A and r_B on the parameters of the experiment (i.e. irrelevance of the locality condition), is not possible. This is because the singlet-state is a quantum mechanical nonfactorizable state. This mathematical nonfactorizability of the spin singlet-state is due to its physical entanglement which has a very important scheme in modern theory of quantum information and computation [8].

Two mathematical points should be explained more about the relations (15) and

(16). The first is that if $n \rightarrow \infty$ then (15) cannot be correct. In fact, in real experiments although n may have large values, never goes to infinity. Here, we appeal to the data of one of real experiments in proton-proton scattering experiment of Laméhi-Rachti and Mittag [6], which exemplifies the above set-up where the total count (n) was 10^4 [7]. The second point is that if θ_{AB}^l s were greater than $\frac{\pi}{2}$, one could not reach (16) from (15). Simply, we can always choose number of runnings n so that the right hand side in (15) be less than $\frac{\pi}{2}$. It is enough only to show that quantum theory is able (is possible) to violate (11).

- 1 J.S. Bell, Physics **1**, No.3, 195 (1964).
- 2 J.F. Clauser and M.A. Horne, Phys. Rev. D **10**, 526 (1974).
- 3 H.P. Stapp, Am.J. Phys. **53**, 306 (1985).
- 4 H.P. Stapp, Found. Phys. **18**, 427 (1988).
- 5 L.E. Ballentine, **Quantum Mechanics**, 440 (Prentice-Hall, 1990).
- 6 M. Laméhi-Rachti and W. Mittig, Phys. Rev. **14**, 2543 (1976).
- 7 Private Communication with Dr. Laméhi-Rachti.
- 8 D. Bouwmeester, et al, **The Physics of Quantum Information**, (Springer, 2000).